

**STATUS OF MINERAL RESOURCE INFORMATION
FOR THE ACOMA INDIAN RESERVATION, NEW MEXICO**

By

C. H. Maxwell
U. S. Geological Survey

M. H. Hibpshman
U. S. Bureau of Mines

Administrative Report BIA-18
1976

CONTENTS

SUMMARY AND CONCLUSIONS	1
INTRODUCTION	1
MINERAL OWNERSHIP	1
GEOLOGY	3
General	3
Sedimentary Rock Units	3
Triassic Rocks	5
Chinle Formation	5
Wingate Sandstone	5
Jurassic Rocks	5
Entrada Sandstone	5
Todilto Limestone	5
Summerville Formation	5
Bluff and Zuni Sandstones	5
Morrison Formation	5
Cretaceous Rocks	6
Dakota Sandstone	6
Mancos Shale	6
Mesaverde Group	6
Tertiary and Quaternary Rocks	7
Gravel Beds	7
Landslides	7
Alluvium	7
Igneous Rocks	7
Mafic Intrusives	7
Volcanic Rocks	8
Structure	8
MINERAL RESOURCES	8
General	8
Energy Resources	8
General	8
Coal	9
Oil and Gas	10
Uranium	10

Nonmetallic Resources	10
General	10
Rock Resources	11
Sand and Gravel	11
Clays	11
Lapidary Materials	11
MARKETS	11
MINERAL LEASING	12
MAP COVERAGE	13
RECOMMENDATIONS	13
REFERENCES	15

SUMMARY AND CONCLUSIONS

Commercial production of mineral resources on the Acoma Indian Reservation has been limited to sand and gravel. Clay is mined by the Indians and used for pottery making, and jet, a form of coal, is collected, polished, and used by Indian silversmiths for settings in jewelry. Rock is quarried and used by tribal members to build homes and other buildings on the reservation. These uses for Indian owned resources will continue and possibly may expand slightly in the future.

Moderate coal resources may exist on the reservation, but the development potential cannot be adequately assessed prior to a detailed field examination. Field work is necessary to: (1) determine the quantity of coal present, (2) locate minable deposits, both strip and underground, (3) obtain samples and ascertain coal quality, and (4) determine if a market exists in the region that would sustain a large mining operation. Uranium of sub-ore grade was discovered on Indian land in 1955, and the deposit might contain ore-grade material because of today's high uranium price. It might be possible to recover the uranium by recently developed in-situ leaching methods.

INTRODUCTION

This report was prepared for the U. S. Bureau of Indian Affairs by the U. S. Geological Survey and the U. S. Bureau of Mines under an agreement to compile and summarize available information on the geology, mineral resources, and potential for mineral development on certain Indian lands.

Source material consisted of published and unpublished reports, and personal communication. There was no field work.

The Acoma Indian Reservation is wholly within Valencia County in northwestern New Mexico ([Figure 1](#)), and is an irregular-shaped area of 245,672 acres (99,497 ha), of which 320 acres (129 ha) consists of allotted lands and 6 acres is held by non-Indians. The tribe also owns five sections of land north of the main reservation area in T. 11 N., Rs. 8 and 9 W.

The reservation, located on the southern margin of the San Juan Basin, is an area of mesas, intervening flat valleys, and spectacular scenery. Interstate Highway 40 paralleled by tracks of the Santa Fe Railway Co. passes through the extreme northern part of the reservation. State and Bureau of Indian Affairs roads provide access to southern and central parts of the area.

Principal towns of the area are Albuquerque (population 243,751), about 60 miles (80.45 km) east, and Grants (population 11,500) about 8 miles northwest of the reservation. The population of Acoma, the main village on the reservation, in 1960, was about 500 (Rand McNally Road Atlas).

MINERAL OWNERSHIP

The Acoma Tribe owns the minerals on most of its land, but in some areas, particularly in the southern part of the reservation, all or some of the minerals are held by others. [Table 1](#) lists areas in which the minerals are not owned by the Indians.

Status of Mineral Resource Information For the Acoma Indian Reservation, New Mexico
C. H. Maxwell and M. H. Hibpshman

TABLE 1
Areas on the Acoma Reservation in Which the Acoma Tribe Does Not Own Mineral Rights

Owner		Section	<u>Location</u>		Minerals Owned
			Township	Range	
New Mexico	Arizona Land Co.	Odd	6 N	8 W	Oil and gas
	Do	Odd	6 N	9 W	Do
	Do	1	6 N	10 W	Do
	Do	3	6 N	10 W	Do
	Do	9	6 N	10 W	Do
	Do	11	6 N	10 W	Do
	Do	13	6 N	10 W	Do
	Do	25	6 N	10 W	Do
	Do	27	6 N	10 W	Do
	Do	35	6 N	10 W	Do
	Do	Odd	7 N	7 W	Do
	Do	Odd	7 N	8 W	Do
	Do	13	7 N	9 W	Do
	Do	23	7 N	9 W	Do
	Do	25	7 N	9 W	Do
	Do	27	8 N	8 W	Do
	Do	35	8 N	8 W	Do
	Do	S½ 25	8 N	8 W	Do
	Do	S½ 27	8 N	8 W	Do
	Do	S½ 29	8 N	8 W	Do
	Do	31	8 N	8 W	Do
	Do	33	8 N	8 W	Do
	Do	35	8 N	8 W	Do
	Do	Odd	8 N	9 W	Do
State of Mexico		32, 36	8 N	8 W	All
Unknown		2	6 N	8 W	Surface minerals
	Do	16	6 N	8 W	Do
	Do	32	6 N	8 W	Do
	Do	36	6 N	8 W	Do
	Do	W½NW¼ 28	6 N	8 W	Do
	Do	W½SW¼ 28	6 N	8 W	Do
	Do	2	8 N	9 W	All
	Do	16	8 N	9 W	Do
	Do	32	8 N	9 W	Do
	Do	36	8 N	9 W	Do
	Do	E½NE¼ 13	8 N	10 W	Oil and gas
U.S. Government		N½ 10	8 N	9 W	All

GEOLOGY

General

The only recent geologic publications of areas within the Acoma reservation are geologic quadrangle maps by Thaden and others (1967), Maxwell (1976) and an open-file map by Knox (1967). Moench (1963a,b, 1964) mapped adjacent areas on the east, and Moench and Schlee (1967) include some discussion pertinent to the Acoma reservation. Geologic maps covering the northern and eastern part of the reservation are in preparation.

The exposed rocks in the region are mostly sandstone and shale with lesser amounts of limestone and coal capped in the southwestern part of the reservation by basaltic lava flows. The sedimentary rocks range in age from about 200 m.y. (million years) to 1 m.y., and are covered over much of the area by surficial material that ranges in age from about 1 m.y. to today's sand and silt. The lava flows are all comparatively recent. The oldest are less than 3 m.y.; samples from Mesa Chivato about 15 miles northwest of the reservation have been dated as 2.6 m.y. (Weber, 1971); the flows in the valley near McCartys are associated with artifacts that indicate an age of less than 1,100 years.

The sedimentary rocks are generally flat lying, but dip gently northwest in part of the area. West of McCartys, in the northwestern corner of the reservation, the rocks dip gently east and southeast.

Sedimentary Rock Units

The distribution of the rock units on the Acoma Reservation and surrounding areas is shown on [Figure 2](#); [Table 2](#) summarizes the stratigraphic sequence and thickness of the rock units. [Figure 3](#) diagrammatically represents the complex inter-tonguing of the Cretaceous rocks in the region.

TABLE 2
Stratigraphic Sequence of Sedimentary Rock Units

Symbols on figure			Approximate	Thickness
2	3		<u>Feet</u>	<u>Metres</u>
		CRETACEOUS		
	Kmv	Mesaverde Group	120	40
	Kpl	Point Lookout Sandstone		
		Crevasse Canyon Formation		
	Kcg	Gibson Coal Member	300	100
	Kcd	Dalton Sandstone Member	120	40
	Km	Mancos Shale		
	Kmm	Mulatto Tongue	250-320	80-100
Kmv		Crevasse Canyon Formation		
	Kcs	Stray Sandstone	20-40	7-13
	Kcdi	Dilco Coal Member	160-180	55-60
	Kg	Gallup Sandstone	50-80	17-27
	Km	Mancos? Shale	100-120	35-40
Kg	Kg	Gallup Sandstone	20-40	7-13
	Km	Mancos? Shale	25-60	8-20
	Kg	Gallup Sandstone	20-30	6-10
	Km	Mancos Shale (main body)	300-350	100-117
	Kd	Dakota Sandstone		
	Kdt	Twowells Sandstone Tongue	0-80	0-27
Kdm	Kmw	Mancos Shale Whitewater Arroyo Tongue	70-140	23-47
	Kdp	Dakota Sandstone, Paguate Tongue	0-50	0-17
	Kmc	Mancos Shale, Clay Mesa Tongue	40-60	13-20
	Kdo	Oak Canyon Member	60-90	20-30
	Kds	basal sandstone units	0-60	0-20
		JURASSIC		
Jm	Jm	Morrison Formation	0-300	0-100
	Jz	Zuni Sandstone	10-300	3-100
	Jb	Bluff Sandstone	150-220	50-70
Ju		Jurassic undivided		
	Js	Summerville Formation	60-150	20-50
	Jtg	Gypsum beds	0-100	0-30
	Jt	Todilto Limestone	0-10	0-3
	Je	Entrada Sandstone	0-300	0-100
		TRIASSIC		
Tru		Triassic rocks undivided	300	100
	Trwr	Wingate Sandstone, Rock Point Member	300	100
	Trc	Chinle Shale	1,000	340

Triassic Rocks

Chinle Formation.--The Chinle Formation is the oldest outcropping sedimentary rock in the area. The Chinle is about 600 feet (200 m) thick in the southeastern part of the reservation; and is variegated red, purple, and gray shale with lenses of mudstone, sandstone, and conglomerate. It is soft and easily eroded and commonly is obscured by landslides; outcrops form a characteristic bedlands topography.

Wingate Sandstone.--The Rock Point Member of the Wingate Sandstone crops out on the mesa slope in the southwestern corner of the reservation. Only a thin wedge is exposed on the reservation but immediately to the south it thickens to about 250 feet (80 m) of light reddish-brown shaly siltstone and silty sandstone and white fine-grained sandstone.

Jurassic Rocks

Entrada Sandstone.--The Entrada Sandstone is composed of a lower light orangish red sandstone member, variable in thickness; a medial silty member, dark red-brown; and an upper eolian crossbedded sandstone, red-brown in the northern part of the area changing to light greenish white toward the southwest. The upper member forms the bold vertical and rounded cliffs along the base of the mesa in the southeastern part of the reservation.

Todilto Limestone.--The Todilto Limestone is a few feet thick east of Acoma Pueblo and pinches

out about 3 miles to the south. It is dark gray thin-bedded to laminated, fetid, petroliferous, silty limestone, and is the host rock for several uranium occurrences east of the Acoma reservation.

Summerville Formation.--The Summerville Formation is siltstone and very fine grained sandstone, red-brown, pink and tan in the northern exposures, changing to light tan and white toward the south.

Bluff and Zuni Sandstones.--The Bluff and Zuni Sandstones are generally mapped together as one unit in the region. The Bluff is a fluvialite sandstone, reddish brown north and east of the reservation, changing to light brown and tan to the west and south. It is thickest in the north, thinning and merging with the Zuni to the south. The Zuni is a light-brown to tan eolian sandstone, thin but variable in thickness in the northern part and thickening to 300 feet or more to the south. The Bluff outcrop is a conspicuous vertical cliff in most exposures, and the Zuni crops out as smoothly rounded steep slopes above the cliff in the northern part of the reservation. To the southwest the two units merge and form spectacular vertical cliffs - several hundred feet high. Acoma Pueblo, Sky City, is at the top of near-vertical cliffs of Bluff and Zuni Sandstone, that extend about 200 feet above the valley floor.

Morrison Formation.--The Morrison Formation is the most important unit in the region, because it is the host rock for all of the important uranium deposits. The Morrison is truncated along an east-west line across the central part of the reservation

by the erosional unconformity at the base of the Dakota Sandstone, which southward cuts down across all of the Jurassic and some of the Triassic rocks.

The Morrison Formation has been divided into four members (Moench and Schlee, 1967). The basal unit is the Recapture Member, followed by the Westwater Canyon Member, the Brush Basin Member, and the Jackpile Sandstone. The Recapture is a grayish red and brownish red sandstone with some interstratified mudstone at the top. It varies from 10 to more than 50 feet thick and is absent locally. It is classified in part as a fossil soil zone (Maxwell, 1976). The next overlying unit is a conglomerate composed of fragments and pebbles, both angular and rounded, of quartz, chert, igneous rocks, pegmatite, quartzite, limestone, siltstone, clay balls, and dinosaur bones, in a matrix of sand, silt, and clay. This conglomerate together with an overlying lenticular sandstone and conglomerate is mapped as the Westwater Canyon Member, and varies from 0 to more than 30 feet thick. It is the major host for the uranium deposits in the Grants-Gallup area. Almost all of the Morrison Formation in the Acoma area is Brushy Basin Member, a green and grayish green mudstone and shale with lenses of sandstone and limestone. The Jackpile Sandstone overlies and locally intertongues with the Brushy Basin. It is a sandstone similar to the Westwater and is the host rock for most of the uranium deposits in the Laguna area, northeast of the Acoma reservation. The distal edge of the Jackpile is exposed near the town of Seama (Figure 2).

Cretaceous Rocks

The Cretaceous rocks in the region comprise a complexly intertongued sequence of marine and continental sediments. Figure 3 diagrammatically represents the intertonguing of the continental Dakota Sandstone and Mesaverde Group with the marine Mancos Shale.

Dakota Sandstone.--In most the Acoma Reservation area the Dakota Sandstone formed from marginal marine and shoreline sedimentation; it becomes progressively more continental toward the south and west. The Dakota is composed of conglomerate and sandstone at the base and is overlain by siltstone with thin shale layers and sandstone lenses, and of sandstone and siltstone tongues interlayered with the Mancos Shale.

Mancos Shale.--The Mancos Shale is dark gray clay and silt shale which weathers to broad flat areas, smooth slopes, and badlands topography, and is locally covered by landslides. It is gradational in some cases into gray-brown siltstone and into intertonguing sandstone.

Mesaverde Group.--The Mesaverde Group is divided into three formations, the Point Lookout Sandstone, the Crevasse Canyon Formation, and the Gallup Sandstone. The group is composed mostly of continental sandstone and siltstone with minor marginal marine siltstone and shale. The only units of potential economic interest in the Acoma area are the Dilco Coal Member and the Gibson Coal Member of the Crevasse Canyon Formation.

The Dilco Coal Member is thin-bedded to lamellar sandstone, siltstone, and shale with numerous interbeds of highly carbonaceous shale and coal, most of which are only a few centimeters thick; the thickest coal bed in outcrop is about 50 cm.

The Gibson Coal Member is present in the northern part of Cebolleta Mesa and on Mesa Negra, beneath slopes covered by landslides and talus. Colluvial material containing chips of very carbonaceous shale and black powdery coalified material is common locally. The Gibson has no coal beds of current commercial importance, but it does have a potential about equal to that of the Dilco.

Tertiary and Quaternary Rocks

Gravel Beds.--The gravel beds shown on the geologic map (Tg) are apparently late Tertiary and early Quaternary in age. The cemented beds at higher elevations are probably correlative with the Santa Fe Formation; the ones at lower elevations in the valleys are probably related to early stages in the erosional development of the present topography. The gravel beds near San Fidel have been used as a source of aggregate and road metal, and the ones near Acomita and 10 miles south of McCartys constitute the only remaining reserves on the reservation.

Landslides.--Landslides are a conspicuous feature of the Acoma reservation. The largest or more conspicuous are shown on the geologic map (Figure 2), others occur on almost every hill and mesa in the region where Chinle shale, Morrison

Formation, or Mancos Shale are capped by sandstone beds or basalt flows. They are more common and larger on the north and west facing slopes, than on the south facing slopes. The landslides, generally in the form of torea blocks (Reiche, 1937) were formed during an early wet period, probably late Pleistocene, followed by progressive changes in climate to the present arid conditions. During the wet period large amounts of water percolated through porous sandstones and basalt, soaked and weakened the underlying shales, and produced numerous landslides. As the volume of precipitation lessened, landslides occurred less frequently and less alluvial material was carried away. Eventually the slides stabilized and the valleys were choked with alluvium which covered many of the lower slides and filled the valleys to form broad flat floors. Flash flooding characteristic of the present arid conditions has produced deep gullies in the alluvium, exhuming some landslide blocks.

Alluvium.--Alluvium, colluvium, and recent wind blown sand cover large areas in the region, and are as much as 100 feet thick in some areas. Only the largest and thickest deposits are shown on the geologic map.

Igneous Rocks

Mafic Intrusives

Diabase dikes and sills generally only a few inches to 1 or 2 feet thick but locally 10 feet or more thick, are common in much of the area. One dike, near Acoma Pueblo and McCartys, has been traced for more than 18 miles. It was quarried near

its juncture with the Crow Point fault and crushed for aggregate on the road from Acomita to Acoma Pueblo.

at the north and south ends. Other faults are similar but have smaller displacements.

Volcanic Rocks

Pyroclastic deposits composed of tuff, ash, lapilli, and bombs, and locally mixed pyroclastics and alluvial gravel occur below and interbedded with the basalt and andesite flows on the flanks of Mount Taylor and on part of Horace Mesa. They range in composition from rhyolitic to basaltic. Volcanic cones occur at several localities on Cebolleta and Horace Mesas. The cones, the source of many of the lava flows, are generally a few tens of feet high but have been considerably eroded so that only massive basalt and coarse fragmental material are left. Late Tertiary or early Quaternary basaltic or andesitic lava flows cap Cebolleta and Horace Mesas and Mesa Negra. Recent basalt flows occur in the valley bottom west of McCartys.

Structure

The Acoma reservation lies within the Acoma Sag, an embayment on the southeastern edge of the San Juan Basin, between the Zuni uplift on the west and the Lucero uplift on the east. The layered rocks are near horizontal with a gentle northwest regional dip. They are slightly deformed locally by small undulations and by numerous faults with small displacements. The Crow Point fault has a displacement of as much as 120 feet in Acoma valley, but diminishes and passes into small folds

MINERAL RESOURCES

General

Mineral resources of the Acoma Reservation have not been studied extensively. The area contains a variety of mineral resources, and may contain undiscovered uranium deposits. Coal occurs within the reservation, but information is sparse. Neither petroleum nor natural gas has been discovered in the area. Sand and gravel, and clay are present, but little development has occurred. Lapidary materials may become important locally. No significant metallic resources are known on the reservation.

Energy Resources

General

Coal is the largest known energy resource on the reservation. Uneconomic uranium resources were discovered at Woods Mesa in 1955, and a radioactive anomaly is known immediately north of the reservation boundary. The possibility of a petroleum and/or natural gas discovery has been rated at 10 percent or less by the Conservation Division of the U. S. Geological Survey.

Coal

Coal occurs in the southernmost part of the reservation and in the five sections of land owned by the tribe in T. 11 N., Rs. 8 and 9 W (Figure 4). The greater resource is probably in the Datil Mountain field to the south, but the area is unexplored and information is sparse. About the only description of the Datil Mountain field available is that of Read, Duffner, Wood, and Zapp (1950) who state:

"The coal in the Datil Mountain field occurs in Upper Cretaceous strata that are correlated with portions of the Mesaverde Formation in the adjacent San Juan River region. The scanty information available concerning the deposits indicates that the coal is of subbituminous rank over most of the area, though locally it may be of bituminous rank. The beds usually are thin and are rarely reported to be more than 3 feet (1 m) in thickness."

"In general, the area is synclinal, but the geologic structure is complicated as a result of some folding, faulting, and igneous intrusion. Until more detailed studies of the coalbeds are made, estimates of the coal reserves must be classed as inferred."

The same authors estimate the total coal resource in the field to be 1.3 billion short tons (1.2 billion mt). However, only about 15 to 20 percent of the field is within reservation boundaries. Therefore, if the 1.3-billion-ton figure is correct,

the resource on the reservation may not exceed 300 million tons (272 million mt). More recent publications on strippable coal reserves of the area--Kottowski, Beaumont, Shomaker (1971) and staff, U. S. Bureau of Mines (1971)--list no strippable reserves for the Datil Mountain field. Similarly, Matson and White (1975) list no underground minable reserves for the Datil Mountain field. However, it should be mentioned that the Datil Mountain field has been little explored and the resources of the field are not accurately known.

Coal in the five tribally owned sections in T. 11 N., Rs. 8 and 9 W., is within the East Mt. Taylor coalfield. It is in the Crevasse Canyon Formation of the Mesaverde Group, which commonly contains two coal-bearing units: the Dilco, or lower coal member, and the Gibson, or upper coal member. Hunt (1936) reports that coal occurs in all of T. 11 N., R. 8 W., and in T. 11 N., R. 9 W., but does not indicate coal in the sections owned by the Indians. However, he did describe coal underlying Indian land on Horace Mesa and reported that the coal does not exceed 14 inches (35.5 cm) thick. Kottowski and Parkhill (1971) reported coalbeds 3 ft (1 m), 5 ft (1.5 m), and 7 ft (2.1 m) thick north of the Indian land in T. 11 N., R. 8 W. A 3-ft. coal bed was mined in the 1930's from two mines about 6 miles (9.6 km) north of the Indian land. Most of the coal in the area is probably in the Gibson Coal Member but the Dilco Coal also is present. Without a detailed field examination, it is impossible to determine if reserves are sufficient to sustain a mining operation.

Oil and Gas

Petroleum and natural gas have not been discovered on Acoma lands, although at least two exploratory wells have been drilled. According to the Conservation Division, U. S. Geological Survey, the nearest petroleum production is probably the Elmer L. White Well No. 3 in sec. 13, T. 15 N., R. 10 W., completed in April 1956, and initially yielding 12 barrels of oil per day. It is probably abandoned now.

The sedimentary sequence is thin, less than 5,000 ft (1,525 m), and the outlook for oil and gas discovery is not promising. The Conservation Division of the Geological Survey rates the oil and gas potential of the reservation as poor, chances of discovery being 10 percent or less.

Uranium

No economic uranium deposits have been discovered, and only one occurrence has been found within the Acoma reservation. Important uranium ore bodies do occur, however, east, west, and north of the reservation in the Laguna, Grants, and Ambrosia Lake districts.

The principal host rocks for uranium in the three districts is in the Jackpile sandstone, and the Westwater Canyon Member of the Morrison Formation. The Todilto Limestone contains small economic uranium deposits east of the reservation.

Only one uranium occurrence has been discovered on the reservation (Figure 5). About 1955, Frost Geophysical Co. explored by drilling an area called Woods Mesa in sec. 5, T. 8 N., R. 7 W., and secs. 20, 29, 32, T. 9 N., R. 7 W. Drilling was

spaced on 1,000-foot centers. Particulars of the drilling program are not available, but reportedly the drill holes penetrated both the Morrison and Todilto Formations and bottomed in the underlying Zuni Sandstone. Low grade uranium mineralization was encountered in both the Morrison and Todilto Formations, but it was not considered economic in 1955.

Indications of radioactive material are within a few hundred feet of the north boundary of the reservation (Figure 5), where two water wells penetrated the Zuni Sandstone. Tests conducted by the U. S. Geological Survey found anomalous concentrations of helium, a product of uranium disintegration, in the top 10 ft. (3 m) of the Zuni aquifer (Maxwell, 1976, oral commun.) The anomaly does not prove the presence of uranium in economic concentrations, but it does indicate a good target area for exploration.

Nonmetallic Resources

General

Nonmetallic resources on the reservation consist of rock, sand and gravel, clays, and lapidary material. These nonmetallic resources are used to a limited extent. Rock is quarried and used on the reservation; clay is mined for making pottery; and jet, a form of coal, is collected for use by Acoma silversmiths. Only sand and gravel have been produced by outside interests.

Rock Resources

Rock resources, such as sandstone, limestone, and basalt, occur on the reservation in vast quantities. Both red and white sandstone, suitable for building and landscaping, abound.

The Acoma people have not made any attempt to produce or market their rock resources except for use in constructing their own living and tribal quarters. It is likely, however, that only a limited local market exists for such products.

Sand and Gravel

Sand and gravel resources on the reservation are not large. Some sand and gravel pits have operated in the past, but none currently are being developed (Figure 6). Sand and gravel on the reservation can be mined by permit; apparently no lease is necessary. Former production was at the common corner of secs. 23, 26, T. 10 N., R. 8 W., by the State Highway Department, whose permit expired in 1968. The State Highway Department also had a permit, that expired in 1967 on 10 acres (4 ha) in the SW $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 25, T. 10 N., R. 8 W., and had permits in sec. 22, and NE $\frac{1}{4}$ NE $\frac{1}{4}$ sec. 21, T. 10 N., R. 8 W., which expired upon completion of Interstate Highway 40. The last royalty figures for which records were available indicate that the Indians received 12 cents per ton for their sand and gravel. Most of the gravel has been exhausted; only minor reserves are left, and these are being conserved for Indian use.

Clays

The Acoma people have used local clay for making pottery. This material is being mined from a deposit in secs. 31 and 32, T. 8 N., R. 7 W. (Figure 6). The raw clay is a dark gray and fires light to medium gray, and occurs as a seam 2"-4" thick, that is overlain by 10 or more feet of sandstone. The quantity of clay present on the reservation is unknown.

Lapidary Materials

Lapidary materials, including agate, jet, chalcedony and petrified wood, occur on the reservation, but with the exception of jet, apparently no attempt has been made to market them. Jet, a dense, hard, black form of coal from an area in the Datil Mountain coalfield (Figure 4) is cut and polished, and set in silver jewelry by Acoma silversmiths.

MARKETS

With the exception of coal, mineral products of the reservation probably will be confined to a local market. If a significant coal resource can be found on Acoma land it might be marketed regionally or nationally.

If economic deposits of uranium are discovered the uranium will be marketed nationally because there is no local market.

MINERAL LEASING

Established Federal regulations require royalty and rental fees for mineral leases on Indian reservations, and, a bond must be posted by the lessee for each lease. Regulations pertaining to mineral leasing may be found in the Code of Federal Regulations, 25 CFR and 30 CFR, Requirements for bonds are as follows:

For less than 80 acres (32.4 ha)	\$1,000
For 80 acres (32.4 ha) and less than 120 acres(48.6 ha)	\$1,500
For 120 acres (48.6 ha) and not more than 160acres (97.2 ha)	\$2,000
For each additional 40 acres (16.2 ha), or part thereof, above 160 acres (97.2 ha)	\$ 500

The bond for minerals other than oil and gas may be less, provided the Secretary of the Interior, with consent of the tribe, agrees; or a lessee may file a bond of \$15,000 for all leases in one State if the total acreage does not exceed 10,240 (4147.2 ha). A lessee may also file a bond of \$75,000 for nationwide coverage.

Lessees may acquire more than one lease, but a single lease may not exceed 2,560 acres (1036.8 ha) for minerals other than coal. Ordinarily, coal leases also are limited to 2,560 acres, but upon application, leases can be combined. Leases may be made for any specified term not to exceed 10 years. A diagram of the procedure for obtaining leases and mining permits is shown in [Figure 7](#).

Rentals for minerals are set at \$1.00 per acre (.4 ha) and a development expenditure of not less than \$10.00 per acre per year.

Unless otherwise authorized by the Commissioner of Indian Affairs, the minimum rates for minerals other than oil and gas shall be as follows:

(a) For substances other than gold, silver, copper, lead, zinc, tungsten, coal asphaltum and allied substances, oil, and gas, the lessee shall pay quarterly or as otherwise provided in the lease, a royalty of not less than 10 percent of the value at the nearest shipping point, of all ores, metals, or minerals marketed.

(b) For gold and silver the lessee shall pay quarterly or as otherwise provided in the lease, a royalty of not less than 10 percent to be computed on the value of bullion as shown by mint returns after deducting forwarding charges to the point of sale; and for copper, lead, zinc, and tungsten, a royalty of not less than 10 percent to be computed on the values of ores and concentrates as shown by

reduction returns after deducting freight charges to the point of sale.

(c) For coal the lessee shall pay quarterly or as otherwise provided in the lease, a royalty of not less than 10 cents per ton of 2,000 pounds (907.2 kg) of mine run, or coal as taken from the mine, including what is commonly called "slack".

In addition to the Federal regulations, the tribe requires that "some" Indians be hired by any firm extracting minerals from reservation lands.

MAP COVERAGE

The reservation area is well covered by various map publications. Topographic coverage is provided by the U. S. Geological Survey topographic quadrangles. The entire area has been mapped on the 7 minute or 1:24,000 scale. Such maps may be purchased from U. S. Geological Survey, Branch of Distribution, Central Region, Box 25286, Denver Federal Center, Denver, Colo. 80225. A list follows of topographic quadrangle maps covering the reservation.

7½-Minute Quadrangles

Acoma Pueblo	Grants SE
Blue Mesa	Laguna Honda
Broom Mountain	Los Pilaes
Cebolleta Peak	McCartys
Crow Point	Mecate Meadow
Cubero	Sand Canyon
East Mesa	

Another source of map coverage for the reservation is the U. S. Bureau of Land Management, which has published both land status master title plats and land status quadrangles. Both series of maps can be ordered from the Bureau of Land Management, Records Section, P. O. Box 1449, Santa Fe, N. Mex. An historical index accompanies the master title plats. The quadrangles, master title plats, and historical indexes should be ordered by township and range.

The New Mexico State Highway Department also publishes useful maps of the reservation. Requests should be addressed to the New Mexico State Highway Department, Duplicating Services, P. O. Box 1149, Santa Fe, N. Mex. 87503. The Highway Department has county road maps available. The State Geological Survey in Socorro also is a good source of map information.

Aerial photographic coverage of the reservation may be purchased from both the Geological Survey and the Department of Agriculture. The agencies within the Department of Agriculture from which photos may be obtained are the U. S. Forest Service and the U. S. Soil Conservation Service. Satellite imagery can be obtained from the U. S. Geological Survey, EROS Data Center, Sioux Falls, S. Dak.

RECOMMENDATIONS

Recommendations for further work on the reservation include:

1. A field examination should be made to determine whether economically minable strip and underground coal reserves exist on the reservation.

It is also recommended that an attempt be made to determine whether a market exists in the region for use of the coal.

2. A field examination including drilling is recommended in the Woods Mesa area and inside the reservation near the two water wells that showed anomalous concentrations of helium to determine whether economic concentrations of uranium exist on the reservation.

3. A field examination should be made to identify the quantity and type of clays present on the reservation.

REFERENCES

- Anderson, E. C., 1954, Occurrences of uranium ores in New Mexico: New Mexico Bur Mines and Mineral Resources Circ. 29, 25 p.
- _____, 1957, The metal resources of New Mexico and their economic features through 1954: New Mexico Bur. Mines and Mineral Resources Bull. 39, 183 p.
- Anderson, R. Y., and Kirkland, D. W., 1960, Origin, varves, and cycles of Jurassic Todilto formation, New Mexico: Am. Assoc. Petroleum Geologists Bull., v. 44, no. 1, p. 37-52.
- Argall, G. O., 1954, Why Anaconda's uranium mines are unique: Mining World, v. 16, no. 10, p. 54-59.
- Ash, S. R., 1959, The Indians of west central New Mexico New Mexico Geol. Soc. 10th Field Conf., west central New Mexico, p. 154-156.
- Baker, A. A., Dane, C. H., and Reeside, J. B., Jr., 1947, Revised correlation of Jurassic formations of parts of Utah, Arizona, New Mexico, and Colorado: Am. Assoc. Petroleum Geologists Bull., v. 31, no. 9, p. 1664-1668.
- Barnes, F. C., and Arnold, E., 1951, Proved and potential oil and gas traps of the San Juan Basin, in New Mexico Geol. Soc. Guidebook 2nd Field Conf., south and west sides of the San Juan Basin, New Mexico and Arizona, 1951, p. 132-140.
- Bates, R. L., 1946, Future oil possibilities of New Mexico: New Mexico Bur. Mines and Mineral Resources Circ. 12, 6 p.
- _____, 1943, Selected bibliography of coal in New Mexico: New Mexico Bur. Mines and Mineral Resources Circ. 8, 3 p.
- _____, 1942, The oil and gas resources of New Mexico, second ed.: New Mexico Bur. Mines and Mineral Resources Bull. 18, 319 p.
- Beaumont, E. C., 1961, Petroleum exploration in part of north-central New Mexico in New Mexico Geol. Soc. Guidebook 12th Field Conf., Albuquerque Country, 1961, p. 175-185.
- Beaumont, E. C., Dane, C. H., and Sears, J. D. 1956, Revised nomenclature of Mesaverde group in San Juan Basin, New Mexico: Am. Assoc. Petroleum Geologists Bull., v. 40. no. 9, p. 2149-2162.
- Benjoucky, T. D., 1947, Contribution of New Mexico's mineral industry to World War II: New Mexico Bur. Mines and Mineral Resources Bull. 27, 81 p.
- Bernick, R. W., 1964, Uranium: Eng. Mining Jour., v. 165, no. 2, p. 116-118, 144.
- _____, 1961, Uranium: Eng. Mining Jour., v. 162, no. 2, p. 88-91.
- Bhappu, R. B., 1961, Economic recovery of selenium by flotation from sandstone ores of New Mexico: New Mexico Bur. Mines and Mineral Resources Circ. 58, 42 p.
- Bieberman, R. A., 1951, Mineral Resources of the San Juan Basin, in New Mexico Geol. Soc. Guidebook 2nd Field Conf., south and west sides of the San Juan Basin, New Mexico and Arizona, 1951, p. 141.
- Burks, M. R., and Schilling, J. H., 1955, Bibliography of New Mexico Geology and Mineral Technology through 1950: New Mexico Bur. Mines and Mineral Resources Bull. 43, 198 p.

- Bryan, Kirk, and McCann, F. T., 1937, The Ceja del Rio Puerco--a border feature of the Basin and Range province in New Mexico, pt. 1, Stratigraphy and structure: Jour. Geology, v. 45, no. 8, p. 801-828.
- Burnham, C. W., Metallogenic Provinces of the United States and northern Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 65, 76 p.
- Campbell, M. R., 1922, The coal fields of the United States general introduction: U. S. Geol. Survey Prof. Paper 100-A, 33 p.
- Cannon, H. L., 1953, Geobotanical reconnaissance near Grants, New Mexico: U. S. Geol. Circ. 264, 8 p.
- Chenoweth, W. L., 1957, Radioactive titaniferous heavy mineral deposits in the San Juan Basin, New Mexico and Colorado, in New Mexico Geol. Soc. Guidebook 8th Field Conf., southwestern San Juan Mountains, Colorado, p. 212-217.
- Clippinger, D. M., 1946, Building blocks from natural lightweight materials of New Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 24, 46 p.
- Colman, H. C., 1959, Origin of uranium ores in the Todilto Limestone, Grants, New Mexico, in New Mexico Geol. Soc. 10th Field Conf., west central New Mexico, p. 159.
- Coleman, R. G., and Delevaux, 1957, Occurrence of selenium in sulfides from some sedimentary rocks of the western United States: Econ. Geol. v. 52, no. 5, p. 499-527.
- Cook, F. S., and Wylie, E. T., 1956, The geology of the Woodrow mine, New Mexico (abs.) Econ. Geol., v. 51, No. 1, p. 112-113.
- Cooper, B. N., and Davenport, H. E., 1965, Travertines and related deposits along the east base of the Lucero uplift, New Mexico: (abs.), GSA Sp. papers, v. 82, p. 296-297.
- Dane, C. H., Bachman, G. O., and Reeside, J. B., Jr., 1957, The Gallup sandstone, its age and stratigraphic relationships south and east of the type locality (N. Mex.), in Four Corners Geol. Soc. Guidebook 2d Field Conf. p. 99-113.
- Darton, N. H., 1920, Gypsum in New Mexico in gypsum deposits of the United States: U. S. Geol. Survey Bull. 697, p. 161-186.
- Devers, V. L., 1962, A history and economic analysis of the uranium industry in New Mexico: M. S. Thesis, Univ. New Mexico, 163 p.
- Dinwiddie, G. A., 1963, Ground water in the vicinity of the Jackpile and Paguate mines, in Geology and Technology of the Grant's uranium region: New Mexico Bur. Mines and Mineral Resources, Mem. 15, p. 217-218.
- Duff, A. I., Jr., and Fritts, J. V., 1951, Land of the San Juan Basin in New Mexico Geol. Soc. Guidebook 2nd Field Conf., south and west sides of the San Juan Basin, New Mexico and Arizona, 1951, p. 161-163.
- Duschatko, R. W., 1953, Fracture studies in the Lucero uplift, New Mexico: U. S. Atomic Energy Comm. RME-3072, Tech. Inf. Service, Oak Ridge, Tenn.
- Ellis, R. W., 1936, Analyses of New Mexico coals: U. S. Bur. Mines Tech. Paper 569, 11 p.
- Ellsworth, P. C., and Mirsky, A., 1952, Preliminary report on relation of structure to uranium mineralization in the Todilto limestone, Grants District, New Mexico: U. S. Atomic Energy Comm. RME 4020, 15 p.

- Elston, W. E., 1967, Summary of the mineral resources of Bernalillo, Sandoval, and Santa Fe Counties, New Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 81, 81 p.
- Engineering and Mining Journal, 1975, New Mexico: v. 176, no. 6, p. 304.
- Fassett, J. E., and Hinds, J. S., 1971, Geology and fuel resources of the Fruitland Formation and Kirtland Shale of the San Juan Basin, New Mexico and Colorado: U. S. Geol. Survey Prof. Paper 676, 76 p.
- File, L. A., 1965, Directory of mines of New Mexico: New Mexico Bur. of Mines and Mineral Resources Circ. 77, 188 p.
- Fischer, R. P., 1956, Uranium-vanadium-copper deposits on the Colorado Plateau, in U. S. Geol. Survey Prof. paper 300, p. 143-154.
- Fischer, R. P. and Stewart, J. H., 1960, Distribution and lithologic characteristics of sandstone beds that contain deposits of copper, vanadium, and uranium: U. S. Geol. Survey Prof. Paper 400 B, p. 42 - B44.
- Fitch, D. C., 1971, Exploration Geology methods in the Grants mineral belt: New Mexico Bur. Mines and Mineral Resources Circ. 118, p. 13-28.
- _____, 1974, Uranium resources of New Mexico: New Mexico Bur. Mines and Mineral Resources Circ. 140, p. 82-107.
- Fitch, R., 1961, Let's look at Lance's uranium mines: Mining World, v. 23, no. 11, p. 23-27.
- Foster, R. W., 1966, Sources for lightweight aggregate in New Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 88, 80 p.
- Foster, R. W., Bieberman, R. A., Chaves, R. R., and Niccum, G. K., 1961, Petroleum developments in New Mexico during 1959: New Mexico Bur. Mines and Mineral Resources Bull. 73, 167 p.
- Foster, R. W., and Grant, P. R., 1974, New Mexico Bureau of Mines and Mineral Resources future of New Mexico's oil and gas resources: Resource Map no. 3,
- Freeman, V. L., and Hilpert, L. S., 1956, Stratigraphy of the Morrison formation in part of northwestern New Mexico: U. S. Geol. Survey Bull. 1030-J, p. 309-334.
- Gableman, J. W., 1956, Uranium deposits in limestone: U. S. Geol. Survey Prof. Paper 300, p. 287-404.
- _____, 1956, Uranium deposits in Paludal black shales, Dakota Sandstone, San Juan Basin, New Mexico: U. S. Geol. Survey Prof. Paper 300, p. 303-319.
- Gadway, K. L., 1959, Cretaceous sediments of the north plains and adjacent areas, McKinley, Valencia, and Catron Counties, New Mexico, in New Mexico Geol. Soc. 10th Field Conf., west central New Mexico, p. 81-84.
- Granger, H. C., 1962, Clays in the Morrison Formation and their spatial relationship to the uranium deposits at Ambrosia Lake, New Mexico: U. S. Geol. Survey Prof. Paper 450D, p. 15-20.
- Granger, H. C., Santos, E. S., Dean, B. G., and Moore, F. B., 1961, Sandstone-type uranium deposits at Ambrosia Lake New Mexico: Econ. Geol. Interim Rept., v. 56, no. 7, p. 1180-1209.
- Graves, J. A., Open pit uranium mining: Mining Eng. AIME, v. 26, no. 8, p. 23-25.

- Gregory, H. E., 1917, Geology of the Navajo Country -- a reconnaissance of parts of Arizona, New Mexico, and Utah: U. S. Geol. Survey Prof. Paper 93, 161 p.
- Griswold, G. B., 1969, Derricks and Mines in Mosaic of New Mexico's Scenery, Rocks and History, ed. P. W. Christianson, and F. E. Kottowski. New Mexico Bur. Mines and Mineral Resources scenic trips to the Geol. past no. 8, p. 145-153.
- Guggione, E., 1974, Field shortages trigger a new uranium rush in New Mexico: Mining Eng., AIME, v. 26, no. 1, p. 16-19.
- Hackman, R. J., 1967, Photogeologic map of the Laguna 2 quadrangle, McKinley, Sandoval, and Valencia Counties, New Mexico: U. S. Geol. Survey Open File Map.
- _____, 1967, Photogeologic map of the NE, NW, and SE quarters of the Laguna 1 quadrangle, Sandoval County, New Mexico: U. S. Geol. Survey open File Map.
- Harshbarger, J. W., Repenning, C. A., and Irwin, J. H., 1957, Stratigraphy of the uppermost Triassic and the Jurassic rocks of the Navajo country: U. S. Geol. Survey Prof. Paper 291, 74 p.
- Hemphill, W. R., 1967, Photogeologic map of the east half of the Laguna 4 quadrangle, Bernalillo, Sandoval, and Valencia Counties, New Mexico: U. S. Geol. Survey Open File Map.
- Herrick, H. N., 1904, Gypsum deposits in New Mexico: U. S. Geol. Survey Bull. 223, p. 89-99.
- Hilpert, L. S., 1955, Resource Appraisal northwestern New Mexico: U. S. Geol. Survey TEI 540, p. 106-107.
- _____, 1961, Structural control of epigenetic uranium deposits in carbonate rocks of northwestern New Mexico: U. S. Geol. Survey Prof. Paper 424B, p. 5-8.
- _____, 1969, Uranium resources of northwestern New Mexico: U. S. Geol. Survey Prof. Paper 602, 166 p.
- Hilpert, L. W., and Freeman, V. L., 1956, Guides to uranium deposits in the Morrison Formation, Gallup-Laguna area, New Mexico: U. S. Geol. Survey Prof. Paper 300, p. 299-302.
- Hunt, C. B., 1936, Geology and fuel resources of the southern part of the San Juan Basin, New Mexico, pt. 2, the Mount Taylor Coal Field: U.S. Geol. Survey Bull. 860B, 1936, 80 p.
- _____, 1938, Igneous geology and structure of the Mount Taylor volcanic field New Mexico: U. S. Geol. Survey Prof. Paper 189B, 80 p.
- Huttl, J. B., 1954, New Mexico uranium: Eng. Mining Jour., v. 155, no. 8, p. 96-99.
- _____, 1956, The Jackpile mine, revisited: Eng. Mining Jour., v. 159, no. 10, p. 104-108.
- Jaster, M. C., 1956, Perlite resources of the United States: U. S. Geol. Survey Bull. 1027-E, 29 p.
- Jensen, M. L., 1958, Sulfur isotopes and the origin of sandstone-type uranium deposits: Econ. Geol. v. 53, no. 5, p. 598-615.
- Jicha, H. L., Jr., 1956, A deposit of high calcium lime rock in Valencia County, New Mexico: New Mexico Bur. Mines and Mineral Resources Circ. 36, 5 p.
- _____, 1958, Geology and mineral resources of Mesa del Oro quadrangle, Socorro and Valencia Counties, New Mexico: N. Mex. Inst. Min. and Technology, State Bur. Mines and Mineral Res. Bull. 56, 67 p.

- Jicha, H. L., and Christina, L. B., 1958, Lexicon of New Mexico geologic names, Precambrian through Paleozoic: New Mexico Bur. Mines and Mineral Res. Bull. 56, 67 p.
- Johnson, D. W., 1902, Notes of a geological reconnaissance in eastern Valencia County, New Mexico: Am. geologist, v. 29, no. 2, p. 80-87.
- _____, 1907, Volcanic necks of the Mount Taylor region, New Mexico: Geol. Soc. of Am. Bull., v. 18, p. 303-324.
- Katayama, N., 1960, Genesis of uranium deposits in sedimentary rocks in Rept. of the 21st session Norden, pt. XV, proceedings of sec. 15, genetic problems of uranium and thorium deposits, Copenhagen, 1960, p. 7-13.
- Kelley, V. C., 1950, Regional structure of the San Juan basin: New Mexico Geol. Soc. Guidebook, First Field Conf., San Juan Basin 1950, p. 101-108.
- _____, 1955, Regional tectonics of the Colorado Plateau and relationship to the origin and distribution of uranium: New Mexico Univ. Geology Pub. 5, 120 p.
- _____, 1963, Chm., Geology and technology of the Grants uranium belt: New Mexico Bur. Mines and Mineral Resources, Mem. 15, 277 p.
- Kelley, V. C., Kittel, D. F., and Melancon, P. E., 1968, Uranium deposits of the Grants region, in Ore deposits of the United States, 1933-1957, ed. John D. Ridge, v. 1, pub. AIME, New York, p. 753-769.
- Kelley, V. C., and Wood, G. H., 1946, Geology of the Lucero uplift, Valencia, Socorro, and Bernalillo Counties, New Mexico: U. S. Geol. Survey Oil and Gas Inv. Prelim. Map 47.
- Kerr, P. F., 1958, Uranium emplacement in the Colorado Plateau: Geol. Soc. of Am. Bull, v. 69, p. 1075-1112.
- King, V. L., 1952, Hospah oil field, McKinley County, New Mexico, in Geol. Symposium, Four Corners region: Four Corners Geol. Soc., p. 95-97.
- Kittel, D. F., 1963, Geology of the Jackpile mine area in geology and technology of the Grants uranium region: New Mexico Bur. Mines and Mineral Resources, Mem. 15, p. 167-176.
- Knox, A. S., 1967, Photogeologic map of the Grants 4 quadrangle, Valencia County, New Mexico: U. S. Geol. Survey Open File
- Koehn, M. A., and Koehn, H. H., 1973, Bibliography of New Mexico geology and mineral technology, 1966-1970: New Mexico Bur. Mines and Mineral Resources Bull. 99, 288 p.
- Konigsmark, T. A., 1955, Color changes and uranium deposits of the upper Morrison Formation northeast flank of the Zuni uplift, New Mexico: U. S. Atomic Energy Comm. RME 76, pt. 1, 15 p.
- _____, 1958, Uranium deposits in the Morrison formation in the northeast flank of the Zuni uplift exclusive of Ambrosia Lake, New Mexico: U. S. Atomic Energy Comm. RME 115, 34 p.
- Kottowski, F. E., and Foster, R. W., 1969, Exploration for mineral resources: New Mexico Bur. Mines and Mineral Resources Circ. 101, 126 p.
- Kottowski, F. E., 1971, Strippable low-sulfur coal resources of the San Juan Basin in New Mexico and Colorado: New Mexico Bur. Mines and Mineral Resources, Mem. 25, 189 p.

- Landis, E. R., Dane, C. H., and Cobban, W. A., 1973, Stratigraphic terminology of the Dakota Sandstone and Mancos Shale, west-central New Mexico: U. S. Geol. Survey Bull. 1372-J.
- Lapp, R. E., 1975, We may find ourselves short of uranium too: *Fortune Magazine* v. XCII, No. 4, p. 151-157, 194, 196, 199.
- Lasky, S. G., and Woolton, T. P., 1933, The metal resources of New Mexico and their economic features: New Mexico Bur. Mines and Mineral Resources Bull. no. 7, 178 p.
- Lovering, T. G., 1956, Radioactive deposits in New Mexico: U.S. Geol. Survey Bull. 1009L, 75 p.
- Mathewson, D. E., and Buzzalini, 1958, Uranium deposits of the Mount Taylor Rio Puerco area in New Mexico (abs.) in New Mexico Geol. Survey Guidebook 8th Ann. Field Conf., southwestern San Juan Mountains, Colorado, September 5, 6, & 7, p. 249.
- Matson, T. K., and White, D. H., Jr., 1975, Reserve base of coal for underground mining in the western United States: U. S. Bureau Mines, Inf. Circ. 8678, 148 p.
- Maxwell, C. H., 1976, Geologic map of the Acoma Pueblo quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-1298 (in press).
- Megrue, G. H., 1962, Summerville alteration associated with uranium mineralization, Laguna and Grant, New Mexico: Ph. D. Thesis, Columbia Univ., 62 p.
- Miesch, A. T., Shoemaker, E. M., Newman, W. L., and Finch, W. I., 1960, Chemical composition as a guide to the size of sandstone-type uranium deposits in Morrison Formation on the Colorado plateau: U. S. Geol. Survey Bull. 1112B, 61 p.
- Mineral and Water Resources of New Mexico, 1965: U. S. Geol. Survey, New Mexico Bur. Mines, and Mineral Industries, prepared for U. S. Senate Comm. on Interior and Insular Affairs, Washington, D. C. 437 p.
- Mining Engineering, 1974, Mineralization and geology of "The uranium capital of the world," AIME, v. 26, no. 8, p. 20-22.
- Mirsky, A., 1955, Jurassic rocks of the Lucero uplift, northwestern New Mexico: M. S. Thesis, Univ. of Arizona 99 p.
- _____, 1953, Preliminary report on uranium mineralization in the Dakota Sandstone Zuni uplift, New Mexico: U. S. Atomic Energy Comm. RME 47, 21 p.
- Moench, R. H., 1962, Properties and paragenesis of coffinite from the Woodrow mine, New Mexico: *Am. Mineralogist*, v. 47, nos. 1 and 2, p. 26-33.
- _____, 1962, Vanadium-rich garnet from Laguna, New Mexico: U. S. Geol. Survey Prof. Paper 450B, p. 67-69.
- _____, 1963a, Geologic map of the Seboyeta quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad Map GQ-207.
- _____, 1963b, Geologic map of the Laguna quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-208.

- _____, 1964a, Geologic map of the Dough Mountain quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-354.
- _____, 1964b, Geology of the South Butte quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-355.
- Moench, R. H., and Puffett, W. P., 1963a, Geologic map of the Arch Mesa quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad Map GQ-211.
- _____, 1963b, Geologic map of the Mesa Gigante quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-212.
- Moench, R. H., and Schlee, J. S., 1967, Geology and Uranium Deposits of the Laguna District, New Mexico: U. S. Geol. Survey Prof. Paper 519, 117 p.
- Moench, R. H., Schlee, J. S., and Bryan, W. B., 1965, Geologic map of the La Gotera quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-371.
- New Mexico Bureau of Mines and Mineral Resources, 1958, New Mexico Energy Resources Map. 1972, Geothermal resources of New Mexico: Resource Map No. 1.
- _____, 1964, Mosaic of New Mexico scenery, rocks, and history: Scenic trip to the geologic past, no. 8, p. 146-154.
- _____, 1958, New Mexico land ownership status map.
- _____, 1958, New Mexico metal resources map.
- _____, 1958, New Mexico non-metal resources map.
- _____, 1971, Strippable low-sulfur coal resources of the San Juan Basin in New Mexico and Colorado. Ed. by Shoemaker, J. W., Beaumont, E. C. and F. E. Kottowski, 189 p.
- New Mexico Geological Society, 1952, Guidebook 3rd Field Conf., Rio Grande County, central New Mexico, 126 p.
- Ninninger, R. D., Everhart, D. L., Adler, H. H., and Kratchman, J., 1960, The genesis of uranium deposits, in Rept. of the 21st sess. Norden, pt. XV, proceeds of sec. 15, genetic problems of uranium and thorium deposits, Copenhagen, p. 40-50.
- Noble, E. A., 1960, Genesis of uranium belts of the Colorado Plateau: Internat. Geol. Cong. 21st Copenhagen, R pt., pt. XV, Proc. sec. 15, Genetic problems of uranium and thorium deposits, 6 figures, p. 26-39.
- Northrop, S. A., 1959, Minerals of New Mexico: Rev. ed. Univ. of New Mexico, press, 1959, 665 p.
- Page, L. R., 1960, The source of uranium in ore deposits, in Rept. of uranium and thorium deposits, Copenhagen, p. 149-64.
- Payne, A. L., 1959, Geology and uranium deposits of the Colorado Plateau; Ph. D. dissert. Stanford Univ., 245 p.
- Phalen, W. C., 1919, Salt resources of the United States: U. S. Geol. Survey Bull. 669, p. 153.
- Pike, W. S., Jr., 1947, Intertonguing marine and nonmarine Upper Cretaceous deposits of New Mexico, Arizona, and southwestern Colorado: Geol. Soc. America Mem. 24, 103 p.
- Rapaport, I., 1955, General geology of the Grants, New Mexico uranium area. Uranium Inf. Digest, v. 2, no. 8, p. 17-18.

- _____, 1952, Interim Report on the ore deposits of the Grants district, New Mexico: U. S. Atomic Energy Comm. RMO 1031, 19 p.
- Rapaport, Irving, Hadfield, J. P., and Olson, R. H., 1952, Jurassic rocks of the Zuni uplift, New Mexico: U. S. Atomic Energy Comm. RMO-642, Tech. Inf. Service, Oak Ridge, Tenn. 47 p.
- Ray, Teri, 1966, Bibliography of New Mexico geology and Mineral Technology, 1961-1965: New Mexico Bur. Mines and Mineral Resources Bull. 90, 124 p.
- Read, C. B., Duffner, R. T., Wood, G. H., and Zapp, A. D., 1950, Coal resources of New Mexico: U. S. Geol. Survey Circ. 89, 24 p.
- Reiche, Parry, 1937, The Toreva block -- a distinctive landslide type: *Jour. Geology*, v. 45, no. 5, p. 538-548.
- Ripley, E. P., 1952, Compilation of State tax law relating to oil, gas, and mining properties in New Mexico: New Mexico Bur. Mines and Mineral Industries Bull. no. 32, 79 p.
- Rosenzweig, A., 1961, Mineralogical notes on the uranium deposits of the Grants and Laguna districts, in New Mexico Geol. Soc. Guidebook 12th Ann. Field Conf., Albuquerque Country, New Mexico: p. 168-171.
- Rosenzweig, A., Gruner, J. W., and Gardiner, L., 1954, Widespread occurrences and character of uraninite in Triassic and Jurassic sediments of the Colorado Plateau: *Econ. Geol.* v. 49, no. 4, p. 351-361.
- Rothrock, H. E., Johnson, C. H., and Hahn, A. D., 1946, Fluorspar resources of New Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 21, 245 p.
- Schilling, C. F., and Schilling, J. H., 1956, Bibliography of New Mexico geology and mineral technology, 1951 through 1955: New Mexico Bur. Mines and Mineral Resources Bull. 56, 136 p.
- _____, 1961, Bibliography of New Mexico geology and mineral technology, 1956-1960: New Mexico Bur. Mines and Mineral Resources Bull. 74, 124 p.
- Schilling, J. H., 1965, Molybdenum resources of New Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 76, 76 p.
- Schlee, J. S., 1963, Sandstone pipes of the Laguna area, New Mexico: *Jour. Sed. Petrology*, v. 33, no. 1, p. 112-123.
- Schlee, J. S., and Moench, R. H., 1961, Properties and genesis of the "Jackpile" sandstone, Laguna, New Mexico, in Peterson, J. A., and Osmond, J. C., eds., *Geometry of sandstone bodies*: Tulsa, Am. Assoc. Petroleum Geologists, p. 134-150.
- _____, 1963a, Geologic map of the Moquino quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-209.
- _____, 1963b, Geologic map of the Mesita quadrangle, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-210.
- Sears, J. D., 1934, Geology and fuel resources of the southern part of the San Juan Basin, New Mexico, pt. 1, The coal field from Gallup eastward toward Mount Taylor: U. S. Geol. Survey Bull. 860A, 29 p.

- Sears, J. D., Hunt, C. B., and Hendricks, T. A., 1941, Transgressive and regressive Cretaceous deposits in southern San Juan Basin, New Mexico: U. S. Geol. Survey Prof. Paper 193-F, p. 101-119.
- Shaler, M. K., 1907, Gypsum in northwestern New Mexico: U. S. Geol. Survey Bull. 315, p. 296-302.
- Sharp, J. V. A., 1955, The uranium deposits in the Morrison Formation, Church Rock area, Grants district, New Mexico: M. A. Thesis, Columbia Univ., 37 p. 5 pls.
- Shoemaker, J. W., 1974, Coal, the energy crisis, and New Mexico: New Mexico Bur. Mines and Mineral Resources Circ. 140, p. 20-28.
- Silver, Caswell, 1948, Jurassic overlap in western New Mexico: Am. Assoc. Petroleum Geologists Bull., v. 32, no. 1, p. 68-81.
- Smith, C. T., 1951, Guidebook of the south and west sides of the San Juan Basin, New Mexico and Arizona: New Mexico 2nd Field Conf.
- Sorensen, E. F. Stotelmeyer, R. B., and Baker, D. H., Jr., 1973, Mineral resources and water requirements for New Mexico mineral industries: New Mexico Bur. Mines and Mineral Resources, Circ. 138, 26 p.
- Stark, J. T., 1956, Geology of the south Manzano Mountains, New Mexico: New Mexico Bur. Mines and Mineral Resources Bull. 34, 46 p.
- Summers, W. K., 1968, Geothermics, New Mexico's untapped resources: New Mexico Bur. Mines and Mineral Resources Circ. 98, 8 p.
- Talmage, S. B., Wooton, T. P., 1937, The non-metallic mineral resources of New Mexico and their economic features: New Mexico Bur. Mines and Mineral Resources Bull. 12, 159 p.
- Thaden, R. E., Merrin, Seymour, and Raup, O. B., 1967, Geologic map of the Grants SE quadrangle, Valencia County, New Mexico: U. S. Geol. Survey Geol. Quad. Map GQ-682.
- Titus, F. B., Jr. 1963, Geology and ground water conditions in eastern Valencia County, New Mexico: New Mexico Bur. Mines and Mineral Resources Ground Water Rept. no. 7, 113 p.
- Towle, C. C. and Rapaport, I., 1952, Uranium deposits of the Grants district, New Mexico: Mining Eng., v. 4, no. 11, p. 1037-1040.
- Truesdell, A. H. and Weeks, A. D., 1960, Paragenesis of uranium ores in Todilto Limestone near Grants, New Mexico: U. S. Geol. Survey Prof. Paper 400 B, p. B-53-B54.
- U. S. Atomic Energy Commission, Guidebook to uranium deposits of the western United States: Open File Rept. 359 p.
- U. S. Bureau of Competition, 1975, Report to the Federal Trade Commission on mineral leasing on Indian lands 226 p.
- U. S. Bureau of Mines, 1971, Strippable reserves of bituminous coal and lignite in the United States: U. S. Bur. Mines Inf. Circ. 8531, 148 p.
- U. S. Code of Federal Regulations, 1974, Title 25 Indians, pt. 171, Indians leasing of tribal lands for mining: Sp. ed. Fed. Register, p. 229-284.
- U. S. Department of Commerce, 1974, Federal and State Indian reservations and Indian trust areas: p. 359-361.
- _____, 1970, U. S. Census of Population for New Mexico: p. 33-11.
- U. S. Geological Survey, 1946, Lucero Uplift, Valenica, Socorro, and Bernallillo Counties, New Mexico: Oil and Gas Inv. Prelim. Map 47.

- Weber, R. H., 1971, K/Ar ages of Tertiary igneous rocks in central and western New Mexico: *Isochron West*, no. 71-1, p. 40-41.
- Weber, R. H., and Kottowski, F. E., 1959, Gypsum resources of New Mexico: *New Mexico Bur. Mines and Mineral Resources Bull.* 68, 68 p.
- Wells, E. H., and Wooton, T. P., 1946, Gold mining and gold deposits in New Mexico: *New Mexico Bur. Mines and Mineral Resources Circ.* 5, 23 p.
- Wengerd, S. A., 1959, Regional geology as related to the petroleum potential of the Lucero region, west central New Mexico, in *New Mexico Geol. Soc. 10th Ann. Field Conf.*, west central New Mexico, p. 121-134.
- White, L., 1975, In-situ leaching opens new uranium reserves in Texas: *Eng. and Min. J.*, v. 176, no. 7, p. 73-80.
- Williams, F. E., Fillo, P. V., and Bloom, P. A., 1964, Barite deposits of New Mexico: *New Mexico Bur. Mines and Mineral Resources Circ.* 76, 46 p.
- Winchester, D. E., 1933, The oil and gas resources of New Mexico: *New Mexico Bur. Mines and Mineral Resources Bull.* 9, 223 p.
- Wright, H. E., Jr., 1946, Tertiary and Quaternary geology of the Lower Rio Puerco area, New Mexico: *Geol. Soc. America Bull.*, v. 57, no. 5, p. 383-456.
- Wooton, T. P., 1930, Geologic literature of New Mexico: *New Mexico Bur. Mines and Mineral Resources Bull.* 5, 127 p.
- Young, W. E. and Delicate, D. T., 1965, Mining methods and costs at Section 23 uranium mine, Homestake-Sapin partners, McKinley County, New Mexico: *U. S. Bur. Mines IC* 8280, 48 p.
- Young, W. H., and Anderson, R. L., 1947, Thickness of bituminous coal and lignite seams mined in the United States in 1945: *Bur. Mines IC* 7442, 17 p.
- Youngblood, I. and Koehn, H. H., 1970, Thesis and dissertations the New Mexico Institute of Mining and Technology: *New Mexico Bur. Mines and Mineral Technol. Circ.* 107, 20 p.




Figure 1. Index map showing infrastructure of Acoma Indian Reservation, New Mexico.




Figure 2. Geologic map of the Acoma Indian Reservation and vicinity (compiled from unpublished maps by C.H. Maxwell, 1976).




Figure 3. Diagrammatic representation of stratigraphic relationships of rock units on and near the Acoma Reservation.




Figure 4. Map showing coal occurrences on the Acoma Indian Reservation.




Figure 5. Map showing uranium occurrences on Acoma Indian Reservation.




Figure 6. Map showing former sand and gravel producing area and clay occurrence on the Acoma Indian Reservation.




Figure 7. Diagram outlining procedures for obtaining leases and mining permits on the Acoma Indian Reservation.